



Mass loss from an ice-sheet drainage basin in West Greenland

Andersen, Morten L.; Bech Andersen, Signe; Stenseng, Lars; Skourup, Henriette; Colgan, William; Kristensen, Steen Savstrup; Merryman Boncori, John Peter; Ahlstrøm, Andreas P.; Fettweis, Xavier; Forsberg, René

Total number of authors:
13

Published in:
Review of Survey activities 2013

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Andersen, M. L., Bech Andersen, S., Stenseng, L., Skourup, H., Colgan, W., Kristensen, S. S., Merryman Boncori, J. P., Ahlstrøm, A. P., Fettweis, X., Forsberg, R., Citterio, M., Box, J. E., & Fausto, R. S. (2014). Mass loss from an ice-sheet drainage basin in West Greenland. In O. Bennike, A. A. Garde, & W. S. Watt (Eds.), *Review of Survey activities 2013* (pp. 87-90). Geological Survey of Denmark and Greenland, GEUS. Geological Survey of Denmark and Greenland (GEUS) Bulletin Vol. 2014 No. 31
<http://www.geus.dk/publications/bull/nr31/index-uk.htm>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Mass loss from an ice-sheet drainage basin in West Greenland

Morten L. Andersen, Signe B. Andersen, Lars Stenseng, Henriette Skourup, William Colgan, Steen S. Kristensen, John P. Merryman Boncori, Andreas P. Ahlstrøm, Xavier Fettweis, René Forsberg, Michele Citterio, Jason E. Box, Dirk van As and Robert S. Fausto

The Greenland ice sheet is losing mass to the ocean at an increasing rate (Thomas *et al.* 2006). During the 1980s the ice sheet was believed to be in near-equilibrium (van den Broeke *et al.* 2009). Within the first decade of the 21st century, however, a net negative balance was observed. Greenland's present rate of ice loss is *c.* 250 Gt yr⁻¹, equivalent to a sea-level rise contribution of *c.* 0.69 mm yr⁻¹. The rate of ice loss has increased over the post 1992 observation period (Shepherd *et al.* 2012).

The ice-sheet mass budget can be partitioned into two main components: (1) surface mass balance (SMB; the net difference between accumulation and surface ablation) and (2) marine ice loss (*D*; iceberg discharge via glacier dynamics plus subsurface melt at the glacier terminus). Over the past decade, the surface mass-balance proportion has accelerated relative to the *D* component, changing from *c.* 50% in 2000–2008 (van den Broeke *et al.* 2009) to more than two thirds (68%) in 2009–2012 (Enderlin *et al.* 2014).

Whereas modern climate models appear to capture the surface mass-balance response to climate change, the physical processes driving variability in glacier discharge are more complex. Recent increases in *D* may be due to: (1) changing force-balance at the ice-ocean interface as suggested by model simulations (e.g. Nick *et al.* 2009, 2013), (2) changing basal lubrication at the ice-bed interface due to increased meltwater availability (Zwally *et al.* 2002; Andersen *et al.* 2010), and/or (3) decreasing ice viscosity due to increasing ice temperature (van der Veen *et al.* 2011). The high spatial variability in these forcing mechanisms and a large sensitivity to local fjord geometry (Nick *et al.* 2013) require basin scale studies of glacier dynamics to elucidate local causes of glacier acceleration.

In 2007 the Programme for Monitoring the Greenland Ice Sheet (PROMICE) was initiated to gain insight into the changing mass balance of the Greenland ice sheet using quantitative meteorological observations, as well as airborne surveys of ice thickness and flow-velocity observations (Ahlstrøm *et al.* 2008). Here we present the first calculations of ice discharge using PROMICE observations, with focus on a West Greenland ice-sheet drainage basin previously defined as 'Basin 7' (Zwally *et al.* 2012; Fig. 1). The *c.* 400 km

long ice-sheet margin within Basin 7 includes the 6 km wide Jakobshavn Isbræ, and several other marine-terminating outlet glaciers, such as Store Gletscher and Rink Isbræ (Figs 1, 2). We combine satellite-derived, ice-surface velocities, airborne ice-thickness measurements, and modelled surface mass balance to assess the dynamic discharge from Basin 7.




Fig. 1. Map of Greenland showing interpolated flight lines for 2007 and 2011, area of computed surface velocities for this study (Fig. 3) and the studied catchment area.




Fig. 2. South-to-north section of the studied basin along the 2007 and 2011 flight lines. For location see Fig. 1.

Data and methods

We estimate the solid ice discharge (D) into the ocean according to the input–output method of Rignot & Kanagaratnam (2006). First we quantify the mass flux (F) discharging across a flux gate, upstream of the boundary between the ice sheet and the ocean (the grounding line), defined by the path of the PROMICE airborne ice-thickness surveys conducted in the summers 2007 and 2011 (Fig. 1). The elevation of this flux gate is $c.$ 1500 m a.s.l. in Basin 7 (Figs 2, 3). The ice-surface and bed elevations determined by the airborne surveys were interpolated to $c.$ 30 m spacing along the flux gate to resolve spatial variability in ice flow.

The flux F at grid point i is computed as $F_i = H_i \cdot L_i \cdot v_i$, where H_i is the ice thickness, L_i is the spacing along the flight line ($c.$ 30 m), and v_i is the depth-averaged ice velocity component that is perpendicular to the flux gate. Ice surface velocities were derived by applying offset tracking to ALOS/PALSAR synthetic aperture radar (SAR) data acquired between November 2009 and February 2010, using the SUSIE processing chain based on the commercial package GAMMA (Merryman Boncori *et al.* 2010; Ahlstrøm *et al.* 2011). Uncertainties associated with the ice velocities were estimated using the method of Mohr & Merryman Boncori (2008) and are under 10% (Fig. 3). We assume a uniform vertical velocity profile, where ice-surface velocity is equivalent to depth-averaged velocity (i.e. ‘plug flow’; Rignot & Kanagaratnam 2006).

With the total basin flux $F (= \sum F_i)$ known, the grounding line discharge (D) can be estimated by adding the spatially integrated surface mass balance (SMB) of the area downstream (‘ds’) of the flux gate: $D = F + \text{SMB}_{\text{ds,ref}}$ where $\text{SMB}_{\text{ds,ref}}$ is a reference period (1961–1990) mean SMB field from the regional climate model MAR v3.2, forced at its boundaries

by ECMWF reanalysis data and run at a spatial resolution of 25 km (Fettweis *et al.* 2013a). Similarly, the mass balance upstream of the flux gate (interior mass balance, IMB) can be computed by subtracting F from the upstream spatially integrated SMB for the reference period: $\text{IMB} = \text{SMB}_{\text{us,ref}} - F$.

Quantification of D allows us to estimate the total mass balance (TMB) of the drainage basin. The TMB value is calculated as $\text{TMB} = \text{SMB}_{\text{tot,yr}} - D$, where $\text{SMB}_{\text{tot,yr}}$ is the yearly SMB spatially integrated across the entire basin.

Estimated uncertainty (σb) on radar-derived bed elevation b values is 80 m and estimated uncertainty (σs) on laser-derived surface elevation observations (s) is 0.1 m. Assuming errors in b and s are random, we take uncertainty in ice thickness (σH) as the sum in quadrature of the fractional uncertainties of σb and σs (e.g. Colgan *et al.* 2008). Uncertainty in flux F at gridpoint i (σF_i) is similarly taken as the sum in quadrature of the fractional uncertainties of σH_i and σv_i , where the latter term is the uncertainty in the annual




Fig. 3. Surface velocities derived from synthetic aperture radar (SAR) data for 2009–2010 used in this study. The contour lines are based on the digital elevation model of the *Greenland ice mapping project* (Howat *et al.* 2014).

depth-averaged velocity at i . We assume no uncertainty in L_i . Uncertainty in the total flux F in the basin is then $\sigma F = \sum \sigma F_i$. We take uncertainty in SMB to be 15% at basin scale (Fettweis *et al.* 2013a) and similarly propagate uncertainties in both F and SMB as the sum in quadrature of fractional uncertainties when assessing the cumulative uncertainty (σD) associated with the grounding-line ice discharge. Uncertainties on TMB and IMB are developed analogously.

The thickness observations were carried out in summer, and we do not account for the difference between summer and winter ice velocities. However, at $c.$ 1500 m a.s.l., we expect the difference between summer and winter ice velocities to be small ($<2\%$; Joughin *et al.* 2008).

Results

Both the upstream flux (F) and downstream discharge (D) in Basin 7 are within the uncertainty of their respective values in 2007 and 2011 (Table 1). Whereas we employ different airborne-derived, ice-geometry data for each year, the velocity field used is identical for the two years (winter 2009/2010 values), as is the surface mass-balance correction (1961–1990 values). The similar mass fluxes indicate that changes in ice geometry along the $c.$ 1500 m contour were slight between 2007 and 2011.

Interior mass-balance values for both years are zero within the uncertainty, which is in good agreement with Zwally *et al.* (2011), who found a slight mass gain of 8 Gt yr^{-1} above 2000 m a.s.l.

The total mass-balance values are also, within uncertainty, similar for 2007 and 2011. Considering the $c.$ 10 Gt yr^{-1} decrease in D , this suggests that yearly fluctuations in the dynamics of major tidewater outlet glaciers in Basin 7 are balanced by variations in surface mass balance. The mean total mass-balance value (-30.5 Gt yr^{-1}) corresponds to a sea-level rise contribution of $c.$ 0.08 mm yr^{-1} , and agrees within uncertainty with a satellite gravimetry-derived total mass-balance estimate of -24 ± 1 Gt yr^{-1} for Basin 7 over the 2004 to 2010 period (Colgan *et al.* 2014), and a 2007 total mass-balance value reported in Rignot *et al.* (2008) of -36.7 Gt yr^{-1} for an analogous West Greenland basin. The 2007 value we present is more negative than a Basin 7 estimate of -14 ± 1 Gt yr^{-1} over the 2003 to 2007 period derived from satellite altimetry (Zwally *et al.* 2011). This latter study, however, preceded the 2007 to 2011 observation period, and may therefore reflect the less negative surface mass-balance regime prior to the observation period (Fettweis *et al.* 2013b).

Table 1. Mass fluxes in Gt per year

Year	Upstream flux (F)	Ice discharge (D)	Interior balance (IMB)	Total mass balance*
2007	79.5 ± 6.1	70.4 ± 6.2	-5.6 ± 12.6	-31.3 ± 8.6
2011	69.7 ± 5.3	60.6 ± 5.5	-4.1 ± 12.3	-29.7 ± 7.2

*Total mass balance = $\text{SMB}_{\text{tot,yr}} - D$

Summary remarks

Rignot & Kanagaratnam (2006) invoked an assumption of negligible changes in ice geometry between their flux gates and the grounding line. As their flux gates are located at the $c.$ 1000 m elevation contour, any dynamic thickening or thinning signals affect a relatively small proportion of the basin area. Given that the PROMICE flux gates are substantially farther inland from the grounding line, we are exploring approaches for explicitly correcting D values for recent changes in ice geometry between the upstream flux gate and the downstream grounding line. This may be particularly relevant in highly dynamic areas, such as the Jakobshavn Isbræ area. A preliminary assessment of such a correction for Basin 7 suggests that the rate of change in downstream ice volume is equivalent to $c.$ 25% of D , which would further decrease the total mass balance by up to 15 Gt yr^{-1} . As more synthetic aperture radar data become available, we will improve the temporal coverage of the PROMICE ice-surface velocity product to annual resolution.

The plug-flow assumption adds a negative bias to the mass-loss estimates by assuming that all flow is caused by sliding at the bed, i.e., the surface speed is equal to the mean flow velocity of the ice column. This may be valid in the fast flowing coastal areas, but higher up on the ice sheet the assumption is less valid, where the surface velocity is a mix of sliding and deformation, and the vertically averaged flow speed can be as low as 80% of the observed surface speed.

In the PROMICE framework, this basin-scale mass-balance assessment will be extended to deliver basin-scale mass-balance and ice-discharge estimates of the entire Greenland ice sheet over multiple observation years. This survey aims to improve portioning of mass loss at basin scale, contributing to improved sea-level rise projections for the Greenland ice sheet.

Acknowledgments

PROMICE is funded by the Danish Ministry of Climate, Energy and Building, and is operated by the Geological Survey of Denmark and Greenland. This paper is contribution number 38 of the Nordic Centre of Excellence SVALI, Stability and Variations of Arctic Land Ice, funded by the Nordic Top-level Research Initiative.

References

- Ahlstrøm, A. & the PROMICE project team 2008: A new programme for monitoring the Greenland ice sheet mass loss. *Geological Survey of Denmark and Greenland Bulletin* **15**, 61–64.
- Ahlstrøm, A. *et al.* 2011: PROMICE 2007–2010. Final report for the establishment phase of the Programme for Monitoring of the Greenland Ice Sheet. *Danmarks og Grønlands Geologiske Undersøgelse Rapport* **2011/118**, 178 pp.
- Andersen, M.L. *et al.* 2010: Spatial and temporal melt variability at Helheim Glacier, East Greenland, and its effect on ice dynamics. *Journal of Geophysical Research, Earth Surface* **115**, F04041.
- Colgan, W., Davis, J. & Sharp, M. 2008: Is the high-elevation region of Devon Ice Cap thickening? *Journal of Glaciology* **54**, 428–436.
- Colgan, W., Abdalati, W., Citterio, M., Csatho, B., Fettweis, X., Luthcke, S., Moholdt, G. & Stober, M. 2014: Hybrid inventory, gravimetry and altimetry (HIGA) mass balance product for Greenland and the Canadian Arctic. *The Cryosphere Discussions* **8**, 537–580.
- Enderlin, E.M., Howat, I.M., Jeong, S., Noh, M.-J., van Angelen, J.H. & van den Broeke, M.R. 2014: An improved mass budget for the Greenland ice sheet. *Geophysical Research Letters* **41**, 866–872.
- Fettweis, X., Franco, B., Tedesco, M., van Angelen, J.H., Lenaerts, J.T.M., van den Broeke, M.R. & Gallee, H. 2013a: Estimating the Greenland ice sheet surface mass balance contribution to future sea level rise using the regional atmospheric climate model MAR. *The Cryosphere* **7**, 469–489.
- Fettweis, X., Hanna, E., Lang, C., Belleflamme, A., Erpicum, M. & Gallée, H. 2013b: Brief communication: Important role of the mid-tropospheric atmospheric circulation in the recent surface melt increase over the Greenland ice sheet. *The Cryosphere* **7**, 241–248.
- Howat, I.M., Negrete, A. & Smith, B. 2014: The Greenland ice mapping project (GIMP) land classification and surface elevation datasets. *The Cryosphere Discussions* **8**, 453–478.
- Merryman Boncori, J.P., Dall, J., Ahlstrøm, A.P. & Andersen, S.B. 2010: Validation and operational measurements with SUSIE – a SAR ice-motion processing chain developed within PROMICE (Programme for the monitoring of the Greenland ice sheet). In: Lacoste-Francis, H. (ed.): *Proceedings of the ESA Living Planet Symposium, Bergen, Norway (ESA SP-686) Noordwijk: ESA Communications ESTEC.*
- Mohr, J.J. & Merryman Boncori, J.P. 2008: An error prediction framework for interferometric SAR data. *IEEE Transactions on Geoscience and Remote Sensing* **46**, 1600–1613.
- Nick, F.M., Vieli, A., Howat, I.M. & Joughin, I. 2009: Large-scale changes in Greenland outlet glacier dynamics triggered at the terminus. *Nature Geoscience* **2**, 110–114.
- Nick, F.M., Vieli, A., Andersen, M.L., Joughin, I., Payne, A.J., Edwards, T.L., Pattyn, F. & van de Wal, R.S.W. 2013: Future sea-level rise from Greenland's main outlet glaciers in a warming climate. *Nature* **497**, 235–238.
- Rignot, E. & Kanagaratnam, P. 2006: Changes in the velocity structure of the Greenland ice sheet. *Science* **311**, 986–990.
- Rignot, E., Box, J.E., Burgess, E. & Hanna, E. 2008: Mass balance of the Greenland ice sheet from 1958 to 2007. *Geophysical Research Letters* **35**, L20502.
- Shepherd, A. *et al.* 2012: A reconciled estimate of ice-sheet mass balance. *Science* **338**, 1183–1189.
- Thomas, R., Frederick, E., Krabill, W., Manizade, S. & Martin, C. 2006: Progressive increase in ice loss from Greenland. *Geophysical Research Letters* **33**, L10503, doi:10.1029/2006GL026075.
- Van den Broeke, M., Bamber, J., Ettema, J., Rignot, E., Schrama, E. & van den Berg, W.J., van Meijgaard, E., Velicogna, I. & Wouters, B. 2009: Partitioning recent Greenland mass loss. *Science* **326**, 984–986.
- Van der Veen, C.J., Plummer, J.C. & Stearns, L.A. 2011: Controls on the recent speed-up of Jakobshavn Isbræ, West Greenland. *Journal of Glaciology* **57**, 770–782.
- Zwally, H., Abdalati, W., Herring, T., Larson, K., Saba, J. & Steffen, K. 2002: Surface melt-induced acceleration of Greenland ice-sheet flow. *Science* **297**, 218–222.
- Zwally, H., Giovinetto, M., Beckley, M. & Saba, J. 2012: Antarctic and Greenland drainage systems, NASA, Goddard Space Flight Center, Cryospheric Sciences Laboratory. http://icesat4.gsfc.nasa.gov/cryo_data/ant_grn_drainage_systems.php.
- Zwally, H. J. *et al.* 2011: Greenland ice sheet mass balance: distribution of increased mass loss with climate warming; 2003–07 versus 1992–2002. *Journal of Glaciology* **57**, 88–102.

Authors' addresses

M.L.A., S.B.A., W.C., A.P.A., M.C., J.E.B., D.v.A. & R.S.F., *Geological Survey of Denmark and Greenland, Øster Voldgade 10, DK-1350 Copenhagen K, Denmark.* E-mail: mola@geus.dk

L.S., H.S., S.S.K. & R.F., *DTU Space, DK-2800 Lyngby, Denmark.*

X.F., *Department of Geography, University of Liège, 4000 Liège, Belgium.*

J.P.M.B., *Istituto Nazionale di Geofisica e Vulcanologia, 00142 Rome, Italy.*